



Cross Reference for Tissue Adhesive		
Tissue Seal –Histoacryl®	Dermabond®	Indermil®
HISTOACRYL® BLUE TOPICAL SKIN ADHESIVE <i>PRODUCT # TS1050044FP</i>	<ul style="list-style-type: none"> •DHV12 •DPP6 •DPPXL6 	IND012
HISTOACRYL® CLEAR TOPICAL SKIN ADHESIVE <i>PRODUCT # TS1050071FP</i>	<i>No Clear Offered Only Violet Available</i>	<i>No Clear Offered Only Yellow Available</i>



GENERAL PRODUCT INFORMATION:

Histoacryl® Topical Skin Adhesive is the proven choice for wound closure with extensive clinical use throughout the United States. Histoacryl® consists of n-Butyl-2 Cyanoacrylate and is twice as strong as the leading tissue adhesive available on the market.

FEATURES AND BENEFITS:

- Histoacryl® requires only one layer of adhesive to achieve optimal bonding strength versus two to three layers required for Dermabond®
- Histoacryl® achieves twice the tensile strength of Dermabond®
- Histoacryl® sets up in 10-30 seconds versus 60-90 seconds per layer for Dermabond®
- Histoacryl® is dispensed via a versatile applicator tip
- Histoacryl® polymerizes on contact with tissue & allows 95%+ of the ampoule to be used due to its non-clogging formulation

PATIENT INSTRUCTIONS

The following information should be shared with the patient:

- If possible, avoid contact with water for the first 24 hours after treatment and minimize contact with water for an additional 7-10 days. Patients may shower or bathe but allow only transient wetting of the treatment site. The site should not be scrubbed, soaked or exposed to prolonged wetness for 7-10 days.
- Keep the wound dry with a non-stick, non-medicinal and water resistant bandage, per your doctor's instructions.
- Do not pull or pick at the wound or bandage.
- Do not apply any medications or cream to the wound.
- Avoid extreme physical activity that might dislodge or impact the wound surface.
- Report any discomfort, redness, drainage, swelling or other concerns regarding your wound to your doctor.

